Jkg=Weight of the meteorograph used including accessories (basket, cords, parachute, and the like), in kilograms. Akm = Distance of the starting place from the landing place

in kilometers.

Ad=Direction from the landing place to the starting place (given in degrees as in (\bar{d})).

(b) Additional observations for part B. Therein belong besides the above-named date under (a) 1 and 2, primarily data on the wind at natural levels.

(c) Cloud observations from such stations which have provided no measurements for group A or B. The data proceed in the same

form as was described under (a) 1 and 2.

(d) Observations at mountain stations. These observations are set off by the symbol M before the name of the station. They give pressure (reduced to standard gravity and 0° C.), temperature, and relative humidity; thereafter data on wind and clouds as explained under (a) 1 and 2.

The following abbreviations are used for the kinds of clouds:

Ci = Cirrus As = Altostratus.St = Stratus.Ns = Nimbostratus. Cs = Cirrostratus.

Ac = Altocumulus. Sc=Stratocumulus. Cb=Cumulonimbus. Ce=Cirrocumulus.

Cu = Cumulus.

The star (*) before the name of the place is used in the meaning

The star (*) before the name of the place is used in the meaning above-mentioned (height instead of geopotential).

There follow now examples of the data in part C.

(a) Dallas 1351 158/2; 1 Ci 315/5, 1 Ac 315 (4200); Akg 0.92; Afa 0.6; Jty Fergusson; Jkg 0.2; Akm 106; Ad 65

Means: Observation at Dallas at 13:51: Wind at the surface from direction 158, velocity 2m/sec; 1/10 Cirrus from direction 315, angular velocity 5, 1/10 altocumulus from direction 315, cloud height 4,200 gdm. The weight of the balloon used was 0.92 kg; the free lift 0.6 kg; the instrument of the Fergusson type weighed 0.2 kg; the starting place was 106 km Fergusson type weighed 0.2 kg; the starting place was 106 km distant from the landing place and lay in direction 65° from

(b) Blue Hill 1240 201-801: 290/14; 801-1524: 300/16; 1524-2582: 290/12; 3200: 280/22; 1 Ac 280/10 (4000), Sc (1150)

Means: Observation at Blue Hill at 12:40: Wind at level from leans: Observation at Blue Hill at 12:40: Wind at level from 201 to 801 gdm from direction 290° at 14 m/sec; in the level from 801 to 1,524 gdm from direction 300° at 16 m/sec; in the level from 1,524 to 2,582 gdm from direction 290° at 12 m/sec; wind in 3,200 gdm from direction 280° at 22 m/sec; 1/10 altocumulus coming from direction 280°, angular velocity 10, height of the altocumulus 4,000 gdm, stratogumulus lia at 1,150 gdm

velocity 10, height of the altocumulus 4,000 gdm, stratocumulus lie at 1,150 gdm.

(c) Aviano 1300 7 Ac -/3, Cu 270 (900)

Means: Observation at Aviano at 13:00: 7/10 altocumulus, direction not known, angular velocity 3, cumulus from direction 270 blowing at 900 gdm height.

(d) M Obir 0600 796.7 74.4 73; 2 Cc 270, Sc 270

Means: Observation at the mountain station Obir at 6:00: pressure: 796.7 mb, temperature 74.4°, relative humidity 73 percent; 2/10 cirrocumulus coming from direction 270°, stratocumulus likewise from 270°. stratocumulus likewise from 270°

In this publication the following symbols are employed: P=Pressure in millibars (mb).

T=Temperature in degrees of the tercentesimal scale (temperature in centesimal degrees plus 270°. minus

U=Relative humidity in percent.

Geopotential in geodynamic meters.

d =Wind direction in 360°.

v = Wind velocity in meters/second (m/sec). G = Time in mean Greenwich time (M. G. T.).Cloud-drift direction in 360°, as wind direction, cloud-drift velocity in angular velocity (1,000 v/h).

Weight in kilograms.

Distance in kilometers

The arrangement of the individual stations proceeds according to regions, as below:

Region A=North and Central America.

Region B=South America.

Region C=Europe, North Africa, Siberia.

Region D=India and East Africa. Region E=West and South Africa. Region F=Australia and Oceania.

Region J=Japan and China.

Within each region, the stations lying farthest north always begin at the same north latitude, the station lying farther west comes

An index of all contributing stations with coordinates makes up the conclusion of the 1932 annual of this publication.

Special comment on the sounding balloon observations made in the United States during this period is in order. Sounding balloon observations were not made on each one of the international days but only on the days as shown in the table below. It should be emphasized that the figures in the third column represent the number of sounding balloons released and not the number of instruments returned.

Day	Number of balloons released	Day	Number of balloons released	Day	Number of bal- loons released
1952 Aug. 10 Aug. 11 Aug. 24 Aug. 25 Sept. 14 Sept. 15 Sept. 17 Sept. 28 Sept. 29 Oct. 5 Oct. 12 Oct. 13 Oct. 26 Oct. 27 Nov. 9 Nov. 10 Nov. 23 Nov. 24 Dec. 14	636361366366366366366366	1932—Contd. Doc. 15. Doc. 28. Doc. 29. 1933 Jan. 11. Jan. 12. Jan. 25. Jan. 26. Feb. 8. Feb. 9. Feb. 22 Feb. 23. Mar. 8. Mar. 9. Mar. 22. Mar. 23. Apr. 12. Apr. 13.	63 6 3 6 3 6 3 6 2 5 3 6	1935—Contd. Apr. 26. Apr. 27. May 10. May 11. May 24. May 25. June 7. June 8. June 21. June 22. July 12. July 13. July 26. July 27. Aug. 9. Aug. 10. Aug. 23. Aug. 24.	363636363636363

In all previous sounding balloon campaigns made in the United States by the Weather Bureau the detailed results of the observations have been published in a Weather Bureau publication. The sounding balloon observations of these 2 years in the United States make an exception as the detailed results are available only in the publication here being reviewed.—Richmond T. Zoch.

BIBLIOGRAPHY

[RICHMOND T. ZOCH, in Charge of Library]

By AMY D. PUTNAM

RECENT ADDITIONS

The following have been selected from among the titles of books recently received as representing those most likely to be useful to Weather Bureau officials in their meteorological work and studies:

Abbot, Charles Greeley.

On the corrections to be applied to silver-disk pyrheliometry. Washington. 1937. (In Smithsonian inst. Smithsonian miscellaneous collections, v. 95, no. 23). Publication 3409. 7 p. 24½ cm.

American national committee. World power conference.

Third World power conference, Washington, D. C., September 7-12, 1936. Washington. 1936. 62 p., tabs. 30 cm.

Bennewitz, Kurt.

Aeronautical instruments. Washington. 1923. 13 p. figs., diagrs. 26½ cm. (Extract from Technische Berichte, Vol. III, No. 5.). [Mimeographed.]

Blair, Thomas A.

Weather elements. A text in elementary meteorology. York. 1937. 401 p. illus., maps, tabs., diagrs. 231/4 cm.

Carroll, Thomas, & McAvoy, Wm. H.

The formation of ice upon exposed parts of an airplane in flight. Washington. 1928. 10 p. pl., diagrs. 26½ cm. (National advisory committee for aeronautics. Technical notes, no. 293 July 1928.) [Mimeographed.]

Cleworth, A. B., & Co., ltd.

Practical humidity tables for engineers, textile manufacturers, etc., based upon Prof. Glaisher's hygrometrical tables. London. 1925. 123 p. tables. 17 cm.

Diseker, Ellis G., & Yoder, Robert E.

Sheet erosion studies on Cecil clay. Auburn, Alabama. 1936. 52 p. illus., tabs., diagrs. 23 cm. (Alabama polytechnic institute. Agricultural experiment station. Bulletin 245, November 1936.)

Federal writers' project.

Washington, city and capital. Federal writers' project, Works progress administration. Washington. 1937. xxvi, 1140 [1] p. incl. front., illus., port., plans. maps (part fold.; part in pocket) 24 cm. (American guide series.)

Franz Josef land, Calm bay. Polar station.

Meteorological observations of the Polar station at Franz Josef Land, Calm Bay, Wintering of 1930-1931. Leningrad. 1933. 51 p. tabls., 32½ cm.

Hersey, M. D.

Aneroid investigations in Germany. Washington. 1921. 9 p. tabs. 26½ cm. [Mimeographed.] (National advisory committee for aeronautics. Technical notes, No. 72 Oct. 1921.)

Kunerth, William, & others.

The amount of luminous and of ultra-violet solar radiation received on certain vertical planes. Ames, Iowa. [1936.] 12 p. figs., tabs. 23 cm.

Luckiesh, Matthew, & Moss, Frank K.

Visibility: its measurement and significance in seeing. Phila., Pa. 1935. p. 431-466, illus., tabs., diagrs. (Repr.: Jnl. Frank. inst. v. 220, no. 4. Oct., 1935.)

Mavis, F. T., & Soucek, Edward.

A summary of hydrologic data, Ralston Creek watershed, 1924–1935. Iowa City, Iowa. 1936. 70 p. illus., fold. maps., tabs., diagrs. 23½ cm. (University of Iowa. Studies in engineering. Bulletin 9.)

The physics of the ionosphere. Menasha, Wisconsin. 1937. 43 p. illus., diagrs. 28 cm. (Reprinted from Reviews of modern physics, Vol. 9, January 1937.)

Stresses produced on an airship flying through gusty air. Washington. 1922. 5 p. 26½ cm. (Mimeographed.) (National advisory committee for aeronautics. Technical notes, No. 111, September, 1922.)

Registrierballonaufstiege in einer tiefen Zyklone. Helsingfors. 1935. 32 p. figs., tabs., diagrs. 24 cm. (Mitteilungen des Meteorologischen instituts der Universität. Helsingfors. No. 26.)

Perret, Frank Alvord.

The eruption of Mt. Pelée, Washington, 1935. 1929–1932. (In Carnegie institution of Washington. Publication no. 458.) 126 p. front., illus., fold. diagr. 29 cm.

Pinkerton, Robert M.

Calibration and lag of a Friez type cup anemometer. Washington. 1930. 8 p. illus., table, diagrs. 26½cm. (National advisory committee for aeronautics, Technical notes, no. 341 June 1930.)

Puppo, Agostino.

Problemi relativi alla definizione del clima solare. 1936. 11 p. 24 cm. (Consiglio nazionale delle richerche. Publicazioni del Comitato per la geodesia e la geofisica, No. 8.)

Elektrotechnik und Witterung. Berlin. 1936. 121 p. tabs., diagrs. 22 cm.

Rittmann, A.

Vulkane und ihre Tätigkeit. Stuttgart. 1936. 188 p. illus., tabs. (part fold.). diagrs. 22 cm.

U. S. Coast and geodetic survey.

Magnetic declination in the United States, 1930, by Daniel L. Hazard. Washington. 1932. 41 p. illus., tabs., fold. map. 23½ cm. (Serial No. 540.)

U. S. Congress. Committee on flood control.

Supplemental report to the Committee on flood control, House of Representatives, Seventy-fourth Congress, Second session on H. R. 12517, to provide for a permanent system of flood control, and for other purposes. Washington. 1936. 70 p. tabs. 23 cm.

U. S. National advisory committee for aeronautics.

Aircraft accidents. Method of analysis. Report prepared by special committee on the nomenclature, subdivision, and classification of aircraft accidents. Washington. 1928. 18 p. tabs., diagr. 28 cm. (National advisory committee for aeronautics. Report No. 308.)

U. S. National resources committee.

Public works planning. National resources committee, December 1936. Washington. 1936. 221 p. maps, tabs., diagrs.

Progress report with statements of coordinating committees. June 15, 1936. Submitted to the President in accordance with Executive order no. 7065, June 7, 1935, by the National resources committee. [Washington. 1936.] 61 p. maps (1 fold.) 28½ cm.

U. S. Weather bureau.

Average conditions of wind and weather, north Atlantic ocean and West Indian waters. Compiled for the information of ocean travelers, from data in the Marine division of the Weather bureau. By W. F. McDonald. Issued May 1936. Washington. 1936. 9 p. 23 cm.

Utterback, C. L., & Jorgensen, Wilhelm.

Scattering of daylight in the sea. Seattle, Washington. 1936. 7 p. figs., tabs. 25½ cm. (Reprint from Journal of Oceanographic society of America, v. 26, June 1936.)

Walker, Gilbert Thomas.

Seasonal weather and its prediction. (From Smithsonian inst. Annual report, 1935. Washington, 1936. p. 117-138, illus. (charts), diagrs. 23 cm.) (Reprinted from the Report of the British association for the advancement of science for 1933.)

Walters, R. C. S.

The nation's water supply. London. 1936. xv, 244 p. illus., maps (part fold. in pocket), tables, diagrs. 23½ cm.

Kuroshio und Golfstrom: eine vergleichende hydrodynamische Untersuchung. Berlin. 1936. 69 p. illus., maps (part fold.), tabs., diagrs. 26 cm. (Veröffentlichungen des Insti-tuts für, Meereskunde an der Universität Berlin. Neue Folge. A. Geograph.-naturwissen. Reihe. Heft 29, Mai 1936.)

SOLAR OBSERVATIONS

NOTE ON THE DETERMINATION OF THE TRANSMISSION OF COLOR SCREENS EMPLOYED IN SOLAR RADIATION INTENSITY MEASUREMENTS FOR THE COMPUTATION OF ATMOSPHERIC TURBIDTY FROM WHICH THE WATER VAPOR CONTENT OF THE ATMOSPHERIC TURBINGS TO TH PHERE IS DETERMINED.

By H. H. Kimball, Research Associate, Harvard University

Angström, and also Feussner, have called attention to the fact that different samples of the Schott-glass filters OG₁ and RG₂ vary somewhat among samples of the same color. The cut-off curve in the shorter wavelength end

of the spectrum is very steep, and the wave lengths between which it occurs vary somewhat with temperature as well as with the individual screens examined. Angström advises that this cut-off curve should be determined for each screen employed, as this would help to determine the transmission of the screen as a whole.

So far as I am aware, this has not generally been done. After Hoelper wrote me that American atmospheric turbidity determinations did not seem to be in accord with European determinations, an attempt was made to determine the transmission of screens that have been